

TABLE 1

Summary of Significant Treatment-Related  
Toxicological Parameters in the Reproductive/  
Developmental and General Toxicity Study of AZT/Clarithromycin

Body Weights

Treatment Regimen	Males	Females-A	Females-B
AZT Alone	no body weight change	decrease in body weight	no body weight change
Clarithromycin Alone	decrease in body weight	decrease in body weight	decrease in gestational body weight
AZT + Clarithromycin	decrease in body weight	decrease in body weight	decrease in gestational body weight

Clinical Pathology

Treatment Regimen	Males	Females-A	Females-B
AZT Alone	-mild decrease in RBC, HGB, and HCT -mild increase in MCV and RDW	-mild decrease in RBC, HGB, and HCT -mild increase in MCV and RDW	-mild increase in MCV and RDW
Clarithromycin Alone	-mild neutrophilia	-neutrophilia -reticulocytopenia -decreased MCV -leukocytosis -increase in BUN and creatinine -increase in ALP, ALT, AST, and bile acids	-reticulocytopenia -decreased MCV
AZT + Clarithromycin	-anemia -increased MCV in lower dose groups -decreased MCV in higher dose groups -reticulocytopenia -thrombocytosis -neutrophilia -lymphopenia -increased RDW -increase in ALP, ALT, AST, and bile acids	-anemia -increased MCV in lower dose groups -decreased MCV in higher dose groups -reticulocytopenia -thrombocytosis -neutrophilia -lymphopenia -increased RDW -increase in BUN and creatinine -increase in ALP, ALT, and bile acids	-increased MCV -increased RDW

TABLE 1 (Continued)

Summary of Significant Treatment-Related  
Toxicological Parameters in the Reproductive/  
Developmental and General Toxicity Study of AZT/Clarithromycin

Histopathology		
Treatment Regimen	Males	Females-A
AZT Alone	Liver -hepatocellular cytoplasmic alteration (slight) Spleen -hematopoietic cell proliferation (slight); hemosiderosis (slight)	Liver -hepatocellular cytoplasmic alteration (slight) Spleen -hematopoietic cell proliferation (slight); hemosiderosis (slight)
Clarithromycin Alone	Liver -cytoplasmic alteration -hepatocyte necrosis -cytoplasmic vacuolization Spleen -red pulp atrophy -lymphoid follicle depletion Bone marrow -depletion Kidney -nephropathy Lymph nodes -depletion Thymus -atrophy Brain, choroid plexus -cytoplasmic vacuolization Heart -cytoplasmic vacuolization of valve -degeneration of atrium -necrosis of myocardium Glandular stomach -cystic degeneration Stomach -fungal infection Salivary gland -necrosis	Liver -cytoplasmic alteration -hepatocyte necrosis -cytoplasmic vacuolization Spleen -red pulp atrophy -lymphoid follicle depletion Bone marrow -depletion Kidney -nephropathy Lymph nodes -depletion Thymus -atrophy Brain, choroid plexus -cytoplasmic vacuolization Heart -cytoplasmic vacuolization of valve -degeneration of atrium -necrosis of myocardium Glandular stomach -cystic degeneration Stomach -fungal infection Salivary gland -necrosis

Note: Histopathological evaluations were performed only on gross lesions in Female-B mice.

TABLE 1 (Continued)

Summary of Significant Treatment-Related  
Toxicological Parameters in the Reproductive/  
Development and General Toxicity Study of AZT/Clarithromycin

## Histopathology (Continued)

Treatment Regimen	Males	Females-A
AZT + Clarithromycin	Liver	Liver
	-cytoplasmic alteration	-cytoplasmic alteration
	-hepatocyte necrosis	-hepatocyte necrosis
	-cytoplasmic vacuolization	-cytoplasmic vacuolization
	Spleen	Spleen
	-lymphoid follicle depletion	-lymphoid follicle depletion
	-red pulp atrophy	-red pulp atrophy
	-hemosiderosis	-hemosiderosis
	Bone marrow	Bone marrow
	-depletion	-depletion
	Kidney	Kidney
	-nephropathy	-nephropathy
	Lymph nodes	Lymph nodes
	-depletion	-depletion
	Thymus	Thymus
	-atrophy	-atrophy
	Brain, choroid plexus	Brain, choroid plexus
	-cytoplasmic vacuolization	-cytoplasmic vacuolization
	Heart	Heart
	-cytoplasmic vacuolization of valve	-cytoplasmic vacuolization of valve
	-degeneration of atrium	-degeneration of atrium
	-necrosis of myocardium	-necrosis of myocardium
	Glandular stomach	Glandular stomach
	-cystic degeneration	-cystic degeneration
	Stomach	Stomach
	-fungal infection	-fungal infection
	Salivary gland	Salivary gland
	-necrosis	-necrosis

Note: Histopathological evaluations were performed only on gross lesions in Female-B mice.

TABLE 1 (Continued)

Summary of Significant Treatment-Related  
Toxicological Parameters in the Reproductive/  
Developmental and General Toxicity Study of AZT/Clarithromycin

Reproductive/Developmental		
Treatment Regimen	Females-A Fetuses	Females-B Pups
AZT Alone	<ul style="list-style-type: none"> <li>-reduced live litter size</li> <li>-increased number of resorptions</li> <li>-slight decline in fetal weight per litter</li> </ul>	<ul style="list-style-type: none"> <li>-no adverse effects</li> </ul>
Clarithromycin Alone	<ul style="list-style-type: none"> <li>-reduced live litter size</li> <li>-no conceptuses delivered in high dose group</li> <li>-slight decline in total weight per litter</li> </ul>	<ul style="list-style-type: none"> <li>-decreased number of litters delivered in high dose groups</li> <li>-decreased litter size</li> <li>-decreased number of pups surviving to post natal day 4</li> <li>-reduced pup weight/litter</li> </ul>
AZT + Clarithromycin	<ul style="list-style-type: none"> <li>-reduced live litter size</li> <li>-increased number of resorptions</li> <li>-no conceptuses delivered in highest combination groups</li> <li>-prominent decline in total weight per litter</li> </ul>	<ul style="list-style-type: none"> <li>-decreased incidence of pregnancy</li> <li>-decreased number of litters delivered in highest dose groups</li> <li>-decreased litter size</li> <li>-decreased number of pups surviving to postnatal day 4</li> <li>-reduced pup weight/litter</li> <li>-increased duration of gestation</li> </ul>